


Algeria

Population 2004 (in thousands)

Age	Male	Female	Total
0-29	10,324	9,915	20,239
30-44	3,376	3,292	6,668
45-59	1,681	1,702	3,383
60-69	564	627	1,191
70-79	317	371	687
80+	73	124	197
Total	16,335	16,030	32,366

Population Pyramid



Highlights:

- The death rate from non-communicable diseases (790 per 100,000 population) was about 3.9 times that for communicable diseases (203 per 100,000). Tobacco was responsible for 10% of all NCDs compared with 6% of all communicable disease related deaths.

- Within the non-communicable disease group, ischaemic heart disease accounted for 112 deaths per 100,000 population aged 30 years and over, with 8% of these deaths attributed to tobacco. Cancer of the trachea, bronchus and lung accounted for 25 deaths per 100,000 population but with 76% of these deaths attributed to tobacco.

- Within the communicable disease group, deaths attributed to tobacco accounted for 10% of all lower respiratory infection deaths and 5% of all tuberculosis deaths.

- The death rate due to tobacco in men was (177 per 100,000 men aged 30 years and over) compared with women (5 per 100,000 women aged 30 years and over). The proportion of deaths attributable to tobacco was almost 17% for men and 0% for women.

- Among those who died prematurely, almost one in every 328 deaths among those aged 30-44 and one in 9 among those aged 45-59 years were attributable to tobacco use. Of those who died in the 45-59 year age bracket, tobacco use accounted for 9.9% of those whose death was due to a lower respiratory infection, 16.3% of deaths due to the category of 'all cardiovascular diseases combined', and 71.5% of those who had died from cancer of the trachea, bronchus and lung.

WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Cause of death	Both sexes					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+	
Death rate: ALL CAUSES	190	637	2,129	5,579	15,653	1,062
Death rate attributable to tobacco	1	70	260	547	835	90
Proportion of deaths attributable to tobacco (%)	0	11	12	10	5	8
Death rate: Communicable diseases	36	104	353	1,043	3,734	203
Death rate attributable to tobacco	0	5	24	70	224	11
Proportion of deaths attributable to tobacco (%)	0	5	7	7	6	6
Death rate: Tuberculosis	3	3	4	4	3	3
Death rate attributable to tobacco	0	0	0	0	0	0
Proportion of deaths attributable to tobacco (%)	0	9	10	10	12	5
Death rate: Lower respiratory infections	8	44	195	604	2,311	107
Death rate attributable to tobacco	0	4	23	69	221	11
Proportion of deaths attributable to tobacco (%)	0	10	12	11	10	10
Death rate: Noncommunicable diseases	111	476	1,676	4,338	11,415	790
Death rate attributable to tobacco	1	65	236	477	612	79
Proportion of deaths attributable to tobacco (%)	0	14	14	11	5	10
Death rate: All malignant neoplasms	18	103	345	789	1,463	141
Death rate attributable to tobacco	0	19	90	184	280	29
Proportion of deaths attributable to tobacco (%)	0	19	26	23	19	21
Death rate: Trachea, bronchus, lung cancers	1	17	78	150	191	25
Death rate attributable to tobacco	0	12	62	120	152	19
Proportion of deaths attributable to tobacco (%)	2	72	79	80	80	76
Death rate: All other malignant neoplasms	17	86	267	639	1,272	116
Death rate attributable to tobacco	0	7	29	64	127	11
Proportion of deaths attributable to tobacco (%)	0	8	11	10	10	9
Death rate: All Cardiovascular diseases	37	193	740	2,126	6,484	373
Death rate attributable to tobacco	0	31	76	118	45	24
Proportion of deaths attributable to tobacco (%)	1	16	10	6	1	6
Death rate: Ischaemic heart disease	7	64	272	670	1,601	112
Death rate attributable to tobacco	0	13	32	43	12	9
Proportion of deaths attributable to tobacco (%)	1	20	12	6	1	8
Death rate: Cerebrovascular disease	15	72	272	778	2,285	136
Death rate attributable to tobacco	0	11	25	36	13	8
Proportion of deaths attributable to tobacco (%)	0	15	9	5	1	6
Death rate: Other cardiovascular diseases	16	57	197	677	2,597	124
Death rate attributable to tobacco	0	8	20	39	20	7
Proportion of deaths attributable to tobacco (%)	1	14	10	6	1	5
Death rate: Respiratory diseases	14	53	197	477	1,052	86
Death rate attributable to tobacco	0	13	64	161	264	23
Proportion of deaths attributable to tobacco (%)	1	24	33	34	25	27
Death rate: Chronic obstructive pulmonary disease	3	21	98	246	481	39
Death rate attributable to tobacco	0	10	53	133	206	19
Proportion of deaths attributable to tobacco (%)	3	48	53	54	43	49
Death rate: Other respiratory diseases	11	33	99	231	572	48
Death rate attributable to tobacco	0	3	12	27	58	5
Proportion of deaths attributable to tobacco (%)	0	9	12	12	10	10

Note 1: The total death rates are crude rates, and not standardized therefore cannot be used for comparison between countries/regions.

Note 2: The estimates above are based on SIR method (see Technical Note), some countries have used other methods which can produce somewhat different but equally valid estimates.

WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Cause of death	Men					Total for ages 30 & over	Women					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+		30-44	45-59	60-69	70-79	80+	
Death rate: ALL CAUSES	197	704	2,461	6,093	15,771	1,052	182	572	1,830	5,139	15,584	1,072
Death rate attributable to tobacco	1	140	549	1,101	2,245	177	-	-	-	75	-	5
Proportion of deaths attributable to tobacco (%)	1	20	22	18	14	17	-	-	-	1	-	0
Death rate: Communicable diseases	47	129	427	1,177	3,799	211	25	80	286	928	3,696	196
Death rate attributable to tobacco	0	10	51	141	601	22	-	-	-	9	-	1
Proportion of deaths attributable to tobacco (%)	0	7	12	12	16	11	-	-	-	1	-	0
Death rate: Tuberculosis	4	5	6	6	6	5	2	2	3	3	2	2
Death rate attributable to tobacco	0	1	1	1	1	0	-	-	-	0	-	0
Proportion of deaths attributable to tobacco (%)	0	12	14	14	18	7	-	-	-	1	-	0
Death rate: Lower respiratory infections	13	54	251	721	2,437	114	2	34	145	504	2,237	101
Death rate attributable to tobacco	0	9	49	139	595	22	-	-	-	9	-	1
Proportion of deaths attributable to tobacco (%)	0	16	20	19	24	19	-	-	-	2	-	1
Death rate: Noncommunicable diseases	100	504	1,916	4,668	11,535	763	124	449	1,460	4,057	11,344	817
Death rate attributable to tobacco	1	131	499	960	1,644	155	-	-	-	66	-	4
Proportion of deaths attributable to tobacco (%)	1	26	26	21	14	20	-	-	-	2	-	0
Death rate: All malignant neoplasms	12	101	413	979	2,054	151	24	105	283	628	1,113	132
Death rate attributable to tobacco	0	39	191	389	751	59	-	-	-	10	-	1
Proportion of deaths attributable to tobacco (%)	1	39	46	40	37	39	-	-	-	2	-	0
Death rate: Trachea, bronchus, lung cancers	2	32	148	285	451	44	1	3	15	35	36	5
Death rate attributable to tobacco	0	25	131	254	409	38	-	-	-	5	-	0
Proportion of deaths attributable to tobacco (%)	2	78	88	89	91	85	-	-	-	15	-	6
Death rate: All other malignant neoplasms	10	69	265	693	1,603	106	24	102	268	592	1,077	126
Death rate attributable to tobacco	0	14	60	135	342	21	-	-	-	4	-	0
Proportion of deaths attributable to tobacco (%)	1	20	23	19	21	20	-	-	-	1	-	0
Death rate: All Cardiovascular diseases	31	199	797	2,071	5,782	327	44	187	689	2,173	6,899	417
Death rate attributable to tobacco	1	63	160	224	120	46	-	-	-	28	-	2
Proportion of deaths attributable to tobacco (%)	2	32	20	11	2	14	-	-	-	1	-	0
Death rate: Ischaemic heart disease	8	83	346	824	1,741	125	6	45	205	539	1,518	100
Death rate attributable to tobacco	0	25	67	86	32	18	-	-	-	6	-	0
Proportion of deaths attributable to tobacco (%)	2	30	19	10	2	15	-	-	-	1	-	0
Death rate: Cerebrovascular disease	12	70	273	685	1,816	110	18	73	270	858	2,564	162
Death rate attributable to tobacco	0	22	52	64	34	15	-	-	-	13	-	1
Proportion of deaths attributable to tobacco (%)	1	31	19	9	2	13	-	-	-	1	-	0
Death rate: Other cardiovascular diseases	12	45	179	562	2,225	93	20	69	214	776	2,817	156
Death rate attributable to tobacco	0	16	42	73	53	13	-	-	-	10	-	1
Proportion of deaths attributable to tobacco (%)	3	35	23	13	2	14	-	-	-	1	-	0
Death rate: Respiratory diseases	13	68	279	651	1,352	103	16	39	124	329	875	70
Death rate attributable to tobacco	0	26	136	319	710	46	-	-	-	25	-	2
Proportion of deaths attributable to tobacco (%)	2	38	49	49	53	44	-	-	-	8	-	2
Death rate: Chronic obstructive pulmonary disease	3	29	152	365	711	52	3	12	50	145	344	26
Death rate attributable to tobacco	0	20	111	264	554	37	-	-	-	22	-	1
Proportion of deaths attributable to tobacco (%)	5	68	73	72	78	70	-	-	-	15	-	5
Death rate: Other respiratory diseases	10	38	126	285	640	51	13	27	75	184	531	44
Death rate attributable to tobacco	0	6	25	55	156	9	-	-	-	3	-	0
Proportion of deaths attributable to tobacco (%)	0	16	20	19	24	17	-	-	-	2	-	0

Note 3: For information on the sources of mortality data, please refer to Annex C, Table C4 of The global burden of disease: 2004 update.