


Guinea

Population 2004 (in thousands)

Age	Male	Female	Total
0-29	3,167	3,051	6,218
30-44	701	678	1,378
45-59	391	404	795
60-69	132	157	289
70-79	55	72	126
80+	10	15	26
Total	4,455	4,377	8,833

Population Pyramid



Highlights:

- The death rate from non-communicable diseases (1126 per 100,000 population) was about 2 times that for communicable diseases (555 per 100,000). Tobacco was responsible for 7% of all NCDs compared with 3% of all communicable disease related deaths.

- Within the non-communicable disease group, ischaemic heart disease accounted for 157 deaths per 100,000 population aged 30 years and over, with 7% of these deaths attributed to tobacco. Cancer of the trachea, bronchus and lung accounted for 12 deaths per 100,000 population but with 56% of these deaths attributed to tobacco.

- Within the communicable disease group, deaths attributed to tobacco accounted for 6% of all lower respiratory infection deaths and 11% of all tuberculosis deaths.

- The death rate due to tobacco in men was (182 per 100,000 men aged 30 years and over) compared with women (11 per 100,000 women aged 30 years and over). The proportion of deaths attributable to tobacco was almost 9% for men and 1% for women.

- Among those who died prematurely, almost one in every 11 deaths among those aged 30-44 and one in 17 among those aged 45-59 years were attributable to tobacco use. Of those who died in the 45-59 year age bracket, tobacco use accounted for 5.4% of those whose death was due to a lower respiratory infection, 10.5% of deaths due to the category of 'all cardiovascular diseases combined', and 53.4% of those who had died from cancer of the trachea, bronchus and lung.

WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Cause of death	Both sexes					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+	
Death rate: ALL CAUSES	722	1,532	3,535	8,137	17,495	1,803
Death rate attributable to tobacco	68	89	128	271	470	95
Proportion of deaths attributable to tobacco (%)	9	6	4	3	3	5
Death rate: Communicable diseases	372	507	859	1,608	3,194	555
Death rate attributable to tobacco	20	12	16	34	88	19
Proportion of deaths attributable to tobacco (%)	5	2	2	2	3	3
Death rate: Tuberculosis	77	75	96	126	141	82
Death rate attributable to tobacco	13	4	3	4	6	9
Proportion of deaths attributable to tobacco (%)	17	5	3	3	4	11
Death rate: Lower respiratory infections	29	149	401	845	1,882	164
Death rate attributable to tobacco	7	8	13	30	81	10
Proportion of deaths attributable to tobacco (%)	24	5	3	4	4	6
Death rate: Noncommunicable diseases	241	909	2,541	6,304	13,958	1,126
Death rate attributable to tobacco	48	76	112	236	382	76
Proportion of deaths attributable to tobacco (%)	20	8	4	4	3	7
Death rate: All malignant neoplasms	38	196	522	1,137	1,928	211
Death rate attributable to tobacco	7	18	32	70	125	17
Proportion of deaths attributable to tobacco (%)	18	9	6	6	6	8
Death rate: Trachea, bronchus, lung cancers	3	12	27	55	76	12
Death rate attributable to tobacco	2	7	15	34	51	7
Proportion of deaths attributable to tobacco (%)	53	53	54	62	67	56
Death rate: All other malignant neoplasms	34	184	495	1,082	1,852	199
Death rate attributable to tobacco	5	11	17	36	74	10
Proportion of deaths attributable to tobacco (%)	15	6	3	3	4	5
Death rate: All Cardiovascular diseases	80	367	1,121	3,108	7,739	504
Death rate attributable to tobacco	31	39	31	47	19	34
Proportion of deaths attributable to tobacco (%)	39	11	3	2	0	7
Death rate: Ischaemic heart disease	15	124	409	969	1,957	157
Death rate attributable to tobacco	7	15	13	18	5	11
Proportion of deaths attributable to tobacco (%)	47	12	3	2	0	7
Death rate: Cerebrovascular disease	32	137	413	1,148	2,738	187
Death rate attributable to tobacco	11	13	10	13	6	12
Proportion of deaths attributable to tobacco (%)	35	10	2	1	0	6
Death rate: Other cardiovascular diseases	32	107	299	991	3,043	160
Death rate attributable to tobacco	13	10	8	15	9	12
Proportion of deaths attributable to tobacco (%)	40	9	3	2	0	7
Death rate: Respiratory diseases	31	103	296	680	1,331	126
Death rate attributable to tobacco	9	19	46	114	226	23
Proportion of deaths attributable to tobacco (%)	29	18	16	17	17	18
Death rate: Chronic obstructive pulmonary disease	6	40	146	349	626	55
Death rate attributable to tobacco	4	15	41	102	196	18
Proportion of deaths attributable to tobacco (%)	69	38	28	29	31	33
Death rate: Other respiratory diseases	25	63	150	332	705	72
Death rate attributable to tobacco	5	4	5	12	30	5
Proportion of deaths attributable to tobacco (%)	19	6	3	4	4	7

Note 1: The total death rates are crude rates, and not standardized therefore cannot be used for comparison between countries/regions.

Note 2: The estimates above are based on SIR method (see Technical Note), some countries have used other methods which can produce somewhat different but equally valid estimates.

WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Cause of death	Men					Total for ages 30 & over	Women					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+		30-44	45-59	60-69	70-79	80+	
Death rate: ALL CAUSES	773	1,791	4,077	8,916	18,926	1,913	669	1,282	3,080	7,543	16,517	1,696
Death rate attributable to tobacco	115	180	280	626	1,158	182	21	-	-	-	-	11
Proportion of deaths attributable to tobacco (%)	15	10	7	7	6	9	3	-	-	-	-	1
Death rate: Communicable diseases	339	613	1,048	1,872	3,693	587	406	405	701	1,408	2,853	523
Death rate attributable to tobacco	37	25	35	80	217	36	3	-	-	-	-	1
Proportion of deaths attributable to tobacco (%)	11	4	3	4	6	6	1	-	-	-	-	0
Death rate: Tuberculosis	115	121	141	189	226	124	39	31	59	78	83	41
Death rate attributable to tobacco	24	8	6	9	15	17	2	-	-	-	-	1
Proportion of deaths attributable to tobacco (%)	21	7	4	5	6	13	5	-	-	-	-	2
Death rate: Lower respiratory infections	46	175	513	1,019	2,213	192	10	124	306	712	1,656	137
Death rate attributable to tobacco	13	16	29	70	201	20	1	-	-	-	-	0
Proportion of deaths attributable to tobacco (%)	28	9	6	7	9	10	8	-	-	-	-	0
Death rate: Noncommunicable diseases	263	998	2,844	6,734	14,861	1,143	218	822	2,287	5,976	13,342	1,110
Death rate attributable to tobacco	78	155	245	546	941	145	18	-	-	-	-	9
Proportion of deaths attributable to tobacco (%)	30	16	9	8	6	13	8	-	-	-	-	1
Death rate: All malignant neoplasms	32	201	613	1,412	2,693	223	43	192	446	927	1,405	200
Death rate attributable to tobacco	12	36	70	162	308	34	2	-	-	-	-	1
Proportion of deaths attributable to tobacco (%)	37	18	11	11	11	15	4	-	-	-	-	0
Death rate: Trachea, bronchus, lung cancers	5	20	50	110	169	20	2	4	8	13	13	4
Death rate attributable to tobacco	3	13	32	79	126	13	1	-	-	-	-	0
Proportion of deaths attributable to tobacco (%)	64	65	65	72	75	67	27	-	-	-	-	7
Death rate: All other malignant neoplasms	28	180	563	1,302	2,524	203	41	188	438	914	1,392	196
Death rate attributable to tobacco	9	23	38	83	181	21	1	-	-	-	-	1
Proportion of deaths attributable to tobacco (%)	32	13	7	6	7	10	3	-	-	-	-	0
Death rate: All Cardiovascular diseases	83	393	1,178	2,991	7,397	471	77	343	1,073	3,197	7,972	536
Death rate attributable to tobacco	50	79	69	108	48	63	12	-	-	-	-	6
Proportion of deaths attributable to tobacco (%)	61	20	6	4	1	13	15	-	-	-	-	1
Death rate: Ischaemic heart disease	20	165	511	1,191	2,245	182	10	84	323	800	1,760	133
Death rate attributable to tobacco	13	31	28	42	13	21	1	-	-	-	-	0
Proportion of deaths attributable to tobacco (%)	66	19	6	3	1	12	9	-	-	-	-	0
Death rate: Cerebrovascular disease	32	139	403	992	2,330	162	33	135	421	1,267	3,017	211
Death rate attributable to tobacco	15	27	22	31	14	20	7	-	-	-	-	4
Proportion of deaths attributable to tobacco (%)	47	20	6	3	1	12	22	-	-	-	-	2
Death rate: Other cardiovascular diseases	30	89	263	808	2,821	128	34	124	329	1,130	3,194	192
Death rate attributable to tobacco	22	20	18	36	21	21	4	-	-	-	-	2
Proportion of deaths attributable to tobacco (%)	71	23	7	4	1	17	11	-	-	-	-	1
Death rate: Respiratory diseases	34	134	416	937	1,750	156	27	72	196	485	1,045	97
Death rate attributable to tobacco	14	38	101	264	557	45	4	-	-	-	-	2
Proportion of deaths attributable to tobacco (%)	41	28	24	28	32	29	15	-	-	-	-	2
Death rate: Chronic obstructive pulmonary disease	8	58	227	526	927	75	5	23	78	213	420	35
Death rate attributable to tobacco	7	31	90	235	483	36	2	-	-	-	-	1
Proportion of deaths attributable to tobacco (%)	81	53	40	45	52	48	46	-	-	-	-	3
Death rate: Other respiratory diseases	26	76	189	411	823	81	23	49	118	272	625	63
Death rate attributable to tobacco	7	7	11	28	75	9	2	-	-	-	-	1
Proportion of deaths attributable to tobacco (%)	28	9	6	7	9	11	8	-	-	-	-	2

Note 3: For information on the sources of mortality data, please refer to Annex C, Table C4 of The global burden of disease: 2004 update.