


Mali

Population 2004 (in thousands)

Age	Male	Female	Total
0-29	4,250	4,261	8,511
30-44	689	786	1,475
45-59	294	407	701
60-69	125	175	301
70-79	81	101	182
80+	42	52	94
Total	5,482	5,783	11,265

Population Pyramid



Highlights:

- The death rate from non-communicable diseases (1771 per 100,000 population) was about 2 times that for communicable diseases (879 per 100,000). Tobacco was responsible for 2% of all NCDs compared with 1% of all communicable disease related deaths.

- Within the non-communicable disease group, ischaemic heart disease accounted for 247 deaths per 100,000 population aged 30 years and over, with 1% of these deaths attributed to tobacco. Cancer of the trachea, bronchus and lung accounted for 9 deaths per 100,000 population but with 51% of these deaths attributed to tobacco.

- Within the communicable disease group, deaths attributed to tobacco accounted for 2% of all lower respiratory infection deaths and 1% of all tuberculosis deaths.

- The death rate due to tobacco in men was (109 per 100,000 men aged 30 years and over) compared with women (0 per 100,000 women aged 30 years and over). The proportion of deaths attributable to tobacco was almost 3% for men and 0% for women.

WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Cause of death	Both sexes					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+	
Death rate: ALL CAUSES	1,075	1,881	4,033	8,960	20,730	2,798
Death rate attributable to tobacco	-	-	117	272	528	49
Proportion of deaths attributable to tobacco (%)	-	-	3	3	3	2
Death rate: Communicable diseases	630	772	1,130	1,760	3,042	879
Death rate attributable to tobacco	-	-	18	33	74	7
Proportion of deaths attributable to tobacco (%)	-	-	2	2	2	1
Death rate: Tuberculosis	125	157	187	250	234	152
Death rate attributable to tobacco	-	-	4	7	8	1
Proportion of deaths attributable to tobacco (%)	-	-	2	3	4	1
Death rate: Lower respiratory infections	68	200	475	846	1,739	255
Death rate attributable to tobacco	-	-	13	26	65	5
Proportion of deaths attributable to tobacco (%)	-	-	3	3	4	2
Death rate: Noncommunicable diseases	305	979	2,761	6,977	17,388	1,771
Death rate attributable to tobacco	-	-	99	239	454	42
Proportion of deaths attributable to tobacco (%)	-	-	4	3	3	2
Death rate: All malignant neoplasms	47	213	564	1,258	2,283	302
Death rate attributable to tobacco	-	-	30	77	162	14
Proportion of deaths attributable to tobacco (%)	-	-	5	6	7	5
Death rate: Trachea, bronchus, lung cancers	0	2	21	53	71	9
Death rate attributable to tobacco	-	-	12	29	42	5
Proportion of deaths attributable to tobacco (%)	-	-	54	54	60	51
Death rate: All other malignant neoplasms	46	211	542	1,205	2,212	293
Death rate attributable to tobacco	-	-	19	49	120	9
Proportion of deaths attributable to tobacco (%)	-	-	3	4	5	3
Death rate: All Cardiovascular diseases	101	396	1,229	3,443	9,787	852
Death rate attributable to tobacco	-	-	27	44	21	7
Proportion of deaths attributable to tobacco (%)	-	-	2	1	0	1
Death rate: Ischaemic heart disease	18	126	440	1,073	2,502	247
Death rate attributable to tobacco	-	-	11	17	6	2
Proportion of deaths attributable to tobacco (%)	-	-	3	2	0	1
Death rate: Cerebrovascular disease	41	150	459	1,278	3,402	311
Death rate attributable to tobacco	-	-	9	12	6	2
Proportion of deaths attributable to tobacco (%)	-	-	2	1	0	1
Death rate: Other cardiovascular diseases	42	120	331	1,093	3,883	294
Death rate attributable to tobacco	-	-	7	14	9	2
Proportion of deaths attributable to tobacco (%)	-	-	2	1	0	1
Death rate: Respiratory diseases	40	107	312	753	1,662	189
Death rate attributable to tobacco	-	-	40	113	259	21
Proportion of deaths attributable to tobacco (%)	-	-	13	15	16	11
Death rate: Chronic obstructive pulmonary disease	8	40	152	387	771	83
Death rate attributable to tobacco	-	-	36	102	226	18
Proportion of deaths attributable to tobacco (%)	-	-	24	26	29	22
Death rate: Other respiratory diseases	32	67	160	366	891	106
Death rate attributable to tobacco	-	-	4	11	33	2
Proportion of deaths attributable to tobacco (%)	-	-	3	3	4	2

Note 1: The total death rates are crude rates, and not standardized therefore cannot be used for comparison between countries/regions.

Note 2: The estimates above are based on SIR method (see Technical Note), some countries have used other methods which can produce somewhat different but equally valid estimates.

WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Mali

Cause of death	Men					Total for ages 30 & over	Women					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+		30-44	45-59	60-69	70-79	80+	
Death rate: ALL CAUSES	1,195	2,293	4,696	9,832	22,949	3,126	971	1,584	3,560	8,262	18,936	2,533
Death rate attributable to tobacco	-	-	280	612	1,181	109	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	-	6	6	5	3	-	-	-	-	-	-
Death rate: Communicable diseases	598	977	1,470	2,143	3,682	984	659	625	888	1,454	2,524	793
Death rate attributable to tobacco	-	-	42	75	166	15	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	-	3	4	5	2	-	-	-	-	-	-
Death rate: Tuberculosis	195	270	296	385	386	243	64	74	108	142	110	79
Death rate attributable to tobacco	-	-	10	15	19	3	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	-	3	4	5	1	-	-	-	-	-	-
Death rate: Lower respiratory infections	89	231	634	1,039	2,098	310	51	177	361	692	1,449	211
Death rate attributable to tobacco	-	-	31	59	146	12	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	-	5	6	7	4	-	-	-	-	-	-
Death rate: Noncommunicable diseases	345	1,083	3,017	7,371	18,905	1,891	271	904	2,577	6,661	16,160	1,675
Death rate attributable to tobacco	-	-	238	537	1,015	94	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	-	8	7	5	5	-	-	-	-	-	-
Death rate: All malignant neoplasms	41	216	650	1,543	3,198	352	52	211	502	1,029	1,542	262
Death rate attributable to tobacco	-	-	73	174	363	31	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	-	11	11	11	9	-	-	-	-	-	-
Death rate: Trachea, bronchus, lung cancers	0	6	45	95	137	17	1	0	4	19	18	3
Death rate attributable to tobacco	-	-	28	64	95	10	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	-	62	67	69	60	-	-	-	-	-	-
Death rate: All other malignant neoplasms	41	210	605	1,448	3,061	335	51	211	497	1,010	1,525	260
Death rate attributable to tobacco	-	-	45	109	268	21	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	-	7	8	9	6	-	-	-	-	-	-
Death rate: All Cardiovascular diseases	108	424	1,253	3,285	9,659	836	95	376	1,212	3,570	9,891	866
Death rate attributable to tobacco	-	-	64	98	47	15	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	-	5	3	0	2	-	-	-	-	-	-
Death rate: Ischaemic heart disease	25	175	545	1,302	2,880	296	12	91	364	890	2,196	207
Death rate attributable to tobacco	-	-	26	37	13	6	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	-	5	3	0	2	-	-	-	-	-	-
Death rate: Cerebrovascular disease	42	152	432	1,099	3,053	281	40	149	478	1,421	3,685	336
Death rate attributable to tobacco	-	-	21	28	13	4	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	-	5	3	0	2	-	-	-	-	-	-
Death rate: Other cardiovascular diseases	40	97	276	884	3,726	259	43	136	370	1,260	4,010	322
Death rate attributable to tobacco	-	-	17	32	21	5	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	-	6	4	1	2	-	-	-	-	-	-
Death rate: Respiratory diseases	45	146	440	1,021	2,184	247	35	79	220	539	1,239	143
Death rate attributable to tobacco	-	-	97	254	580	46	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	-	22	25	27	19	-	-	-	-	-	-
Death rate: Chronic obstructive pulmonary disease	11	62	240	574	1,128	122	6	25	88	238	483	52
Death rate attributable to tobacco	-	-	87	229	506	41	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	-	36	40	45	34	-	-	-	-	-	-
Death rate: Other respiratory diseases	35	84	200	446	1,057	125	29	54	132	301	757	91
Death rate attributable to tobacco	-	-	10	26	74	5	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	-	5	6	7	4	-	-	-	-	-	-

Note 3: For information on the sources of mortality data, please refer to Annex C, Table C4 of The global burden of disease: 2004 update.