


Mozambique

Population 2004 (in thousands)

Age	Male	Female	Total
0-29	7,131	7,214	14,345
30-44	1,368	1,702	3,070
45-59	775	892	1,667
60-69	272	355	627
70-79	126	173	299
80+	27	43	70
Total	9,699	10,379	20,078

Population Pyramid



Highlights:

- The death rate from non-communicable diseases (1062 per 100,000 population) was about 0.9 times that for communicable diseases (1163 per 100,000). Tobacco was responsible for 0% of all NCDs compared with 0% of all communicable disease related deaths.

- Within the non-communicable disease group, ischaemic heart disease accounted for 150 deaths per 100,000 population aged 30 years and over, with 0% of these deaths attributed to tobacco. Cancer of the trachea, bronchus and lung accounted for 4 deaths per 100,000 population but with 2% of these deaths attributed to tobacco.

- Within the communicable disease group, deaths attributed to tobacco accounted for 0% of all lower respiratory infection deaths and 0% of all tuberculosis deaths.

- The death rate due to tobacco in men was (1 per 100,000 men aged 30 years and over) compared with women (1 per 100,000 women aged 30 years and over). The proportion of deaths attributable to tobacco was almost 0% for men and 0% for women.

WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Cause of death	Both sexes					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+	
Death rate: ALL CAUSES	1,521	1,881	3,480	7,738	17,544	2,359
Death rate attributable to tobacco	-	-	-	15	38	1
Proportion of deaths attributable to tobacco (%)	-	-	-	0	0	0
Death rate: Communicable diseases	1,234	967	950	1,519	3,099	1,163
Death rate attributable to tobacco	-	-	-	2	5	0
Proportion of deaths attributable to tobacco (%)	-	-	-	0	0	0
Death rate: Tuberculosis	104	138	180	205	197	129
Death rate attributable to tobacco	-	-	-	0	1	0
Proportion of deaths attributable to tobacco (%)	-	-	-	0	0	0
Death rate: Lower respiratory infections	9	82	296	720	1,781	121
Death rate attributable to tobacco	-	-	-	2	4	0
Proportion of deaths attributable to tobacco (%)	-	-	-	0	0	0
Death rate: Noncommunicable diseases	184	778	2,343	5,943	14,043	1,062
Death rate attributable to tobacco	-	-	-	13	33	1
Proportion of deaths attributable to tobacco (%)	-	-	-	0	0	0
Death rate: All malignant neoplasms	31	169	480	1,068	1,872	197
Death rate attributable to tobacco	-	-	-	2	11	0
Proportion of deaths attributable to tobacco (%)	-	-	-	0	1	0
Death rate: Trachea, bronchus, lung cancers	0	3	12	29	41	4
Death rate attributable to tobacco	-	-	-	1	3	0
Proportion of deaths attributable to tobacco (%)	-	-	-	2	6	2
Death rate: All other malignant neoplasms	30	166	468	1,038	1,831	192
Death rate attributable to tobacco	-	-	-	2	8	0
Proportion of deaths attributable to tobacco (%)	-	-	-	0	0	0
Death rate: All Cardiovascular diseases	62	315	1,038	2,936	7,875	488
Death rate attributable to tobacco	-	-	-	5	1	0
Proportion of deaths attributable to tobacco (%)	-	-	-	0	0	0
Death rate: Ischaemic heart disease	11	106	379	919	1,993	150
Death rate attributable to tobacco	-	-	-	1	0	0
Proportion of deaths attributable to tobacco (%)	-	-	-	0	0	0
Death rate: Cerebrovascular disease	26	119	388	1,098	2,820	182
Death rate attributable to tobacco	-	-	-	2	0	0
Proportion of deaths attributable to tobacco (%)	-	-	-	0	0	0
Death rate: Other cardiovascular diseases	26	90	272	919	3,063	155
Death rate attributable to tobacco	-	-	-	2	0	0
Proportion of deaths attributable to tobacco (%)	-	-	-	0	0	0
Death rate: Respiratory diseases	24	86	270	637	1,316	117
Death rate attributable to tobacco	-	-	-	5	21	1
Proportion of deaths attributable to tobacco (%)	-	-	-	1	2	0
Death rate: Chronic obstructive pulmonary disease	5	34	132	326	610	51
Death rate attributable to tobacco	-	-	-	5	19	0
Proportion of deaths attributable to tobacco (%)	-	-	-	1	3	1
Death rate: Other respiratory diseases	19	53	137	312	706	66
Death rate attributable to tobacco	-	-	-	1	2	0
Proportion of deaths attributable to tobacco (%)	-	-	-	0	0	0

Note 1: The total death rates are crude rates, and not standardized therefore cannot be used for comparison between countries/regions.

Note 2: The estimates above are based on SIR method (see Technical Note), some countries have used other methods which can produce somewhat different but equally valid estimates.

WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Cause of death	Men					Total for ages 30 & over	Women					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+		30-44	45-59	60-69	70-79	80+	
Death rate: ALL CAUSES	1,519	2,281	4,075	8,495	18,757	2,540	1,522	1,533	3,024	7,190	16,793	2,213
Death rate attributable to tobacco	-	-	-	-	100	1	-	-	-	26	-	1
Proportion of deaths attributable to tobacco (%)	-	-	-	-	1	0	-	-	-	0	-	0
Death rate: Communicable diseases	1,179	1,219	1,164	1,772	3,570	1,243	1,278	748	785	1,336	2,807	1,098
Death rate attributable to tobacco	-	-	-	-	12	0	-	-	-	3	-	0
Proportion of deaths attributable to tobacco (%)	-	-	-	-	0	0	-	-	-	0	-	0
Death rate: Tuberculosis	182	236	269	354	372	218	42	53	111	97	89	56
Death rate attributable to tobacco	-	-	-	-	1	0	-	-	-	0	-	0
Proportion of deaths attributable to tobacco (%)	-	-	-	-	0	0	-	-	-	0	-	0
Death rate: Lower respiratory infections	7	72	365	830	2,035	126	11	91	243	641	1,623	116
Death rate attributable to tobacco	-	-	-	-	11	0	-	-	-	3	-	0
Proportion of deaths attributable to tobacco (%)	-	-	-	-	1	0	-	-	-	0	-	0
Death rate: Noncommunicable diseases	163	847	2,640	6,361	14,747	1,087	201	718	2,114	5,641	13,606	1,042
Death rate attributable to tobacco	-	-	-	-	87	1	-	-	-	23	-	1
Proportion of deaths attributable to tobacco (%)	-	-	-	-	1	0	-	-	-	0	-	0
Death rate: All malignant neoplasms	21	171	569	1,334	2,642	216	39	168	412	875	1,395	181
Death rate attributable to tobacco	-	-	-	-	28	0	-	-	-	4	-	0
Proportion of deaths attributable to tobacco (%)	-	-	-	-	1	0	-	-	-	0	-	0
Death rate: Trachea, bronchus, lung cancers	0	4	14	26	49	4	1	2	11	31	36	4
Death rate attributable to tobacco	-	-	-	-	7	0	-	-	-	1	-	0
Proportion of deaths attributable to tobacco (%)	-	-	-	-	14	2	-	-	-	4	-	2
Death rate: All other malignant neoplasms	20	167	556	1,307	2,593	211	38	165	400	844	1,359	177
Death rate attributable to tobacco	-	-	-	-	21	0	-	-	-	3	-	0
Proportion of deaths attributable to tobacco (%)	-	-	-	-	1	0	-	-	-	0	-	0
Death rate: All Cardiovascular diseases	52	333	1,096	2,824	7,373	459	71	299	994	3,017	8,187	511
Death rate attributable to tobacco	-	-	-	-	3	0	-	-	-	9	-	1
Proportion of deaths attributable to tobacco (%)	-	-	-	-	0	0	-	-	-	0	-	0
Death rate: Ischaemic heart disease	13	142	479	1,135	2,257	180	9	74	301	762	1,829	126
Death rate attributable to tobacco	-	-	-	-	1	0	-	-	-	2	-	0
Proportion of deaths attributable to tobacco (%)	-	-	-	-	0	0	-	-	-	0	-	0
Death rate: Cerebrovascular disease	20	119	378	944	2,340	157	30	119	395	1,210	3,117	203
Death rate attributable to tobacco	-	-	-	-	1	0	-	-	-	4	-	0
Proportion of deaths attributable to tobacco (%)	-	-	-	-	0	0	-	-	-	0	-	0
Death rate: Other cardiovascular diseases	18	73	238	745	2,775	122	31	106	298	1,046	3,241	181
Death rate attributable to tobacco	-	-	-	-	1	0	-	-	-	3	-	0
Proportion of deaths attributable to tobacco (%)	-	-	-	-	0	0	-	-	-	0	-	0
Death rate: Respiratory diseases	21	114	385	886	1,731	148	26	63	181	457	1,060	91
Death rate attributable to tobacco	-	-	-	-	55	1	-	-	-	9	-	0
Proportion of deaths attributable to tobacco (%)	-	-	-	-	3	0	-	-	-	2	-	1
Death rate: Chronic obstructive pulmonary disease	5	49	210	497	913	74	4	20	72	201	422	33
Death rate attributable to tobacco	-	-	-	-	51	1	-	-	-	8	-	0
Proportion of deaths attributable to tobacco (%)	-	-	-	-	6	1	-	-	-	4	-	1
Death rate: Other respiratory diseases	16	65	175	388	818	74	22	43	109	256	637	59
Death rate attributable to tobacco	-	-	-	-	4	0	-	-	-	1	-	0
Proportion of deaths attributable to tobacco (%)	-	-	-	-	1	0	-	-	-	0	-	0

Note 3: For information on the sources of mortality data, please refer to Annex C, Table C4 of The global burden of disease: 2004 update.