


Seychelles

Population 2004 (in thousands)

Age	Male	Female	Total
0-29	21	21	42
30-44	10	10	20
45-59	7	7	14
60-69	2	2	4
70-79	1	2	3
80+	0	1	1
Total	42	43	85

Population Pyramid



Highlights:

- The death rate from non-communicable diseases (1077 per 100,000 population) was about 7 times that for communicable diseases (153 per 100,000). Tobacco was responsible for 6% of all NCDs compared with 2% of all communicable disease related deaths.

- Within the non-communicable disease group, ischaemic heart disease accounted for 150 deaths per 100,000 population aged 30 years and over, with 6% of these deaths attributed to tobacco. Cancer of the trachea, bronchus and lung accounted for 17 deaths per 100,000 population but with 69% of these deaths attributed to tobacco.

- Within the communicable disease group, deaths attributed to tobacco accounted for 5% of all lower respiratory infection deaths and 6% of all tuberculosis deaths.

- The death rate due to tobacco in men was (124 per 100,000 men aged 30 years and over) compared with women (23 per 100,000 women aged 30 years and over). The proportion of deaths attributable to tobacco was almost 7% for men and 2% for women.

WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Cause of death	Both sexes					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+	
Death rate: ALL CAUSES	212	796	2,435	5,255	15,746	1,316
Death rate attributable to tobacco	-	32	262	565	-	72
Proportion of deaths attributable to tobacco (%)	-	4	11	11	-	5
Death rate: Communicable diseases	29	86	201	655	2,072	153
Death rate attributable to tobacco	-	1	4	27	-	2
Proportion of deaths attributable to tobacco (%)	-	1	2	4	-	2
Death rate: Tuberculosis	1	-	5	19	-	2
Death rate attributable to tobacco	-	-	-	2	-	0
Proportion of deaths attributable to tobacco (%)	-	-	-	10	-	6
Death rate: Lower respiratory infections	3	25	53	164	792	44
Death rate attributable to tobacco	-	1	4	25	-	2
Proportion of deaths attributable to tobacco (%)	-	4	8	15	-	5
Death rate: Noncommunicable diseases	118	587	2,133	4,547	13,674	1,077
Death rate attributable to tobacco	-	31	258	538	-	70
Proportion of deaths attributable to tobacco (%)	-	5	12	12	-	6
Death rate: All malignant neoplasms	26	184	522	920	756	202
Death rate attributable to tobacco	-	13	134	229	-	32
Proportion of deaths attributable to tobacco (%)	-	7	26	25	-	16
Death rate: Trachea, bronchus, lung cancers	-	9	51	145	-	17
Death rate attributable to tobacco	-	3	39	115	-	12
Proportion of deaths attributable to tobacco (%)	-	29	76	79	-	69
Death rate: All other malignant neoplasms	26	175	470	775	756	185
Death rate attributable to tobacco	-	10	95	114	-	20
Proportion of deaths attributable to tobacco (%)	-	6	20	15	-	11
Death rate: All Cardiovascular diseases	52	238	1,023	2,121	9,422	564
Death rate attributable to tobacco	-	17	98	199	-	28
Proportion of deaths attributable to tobacco (%)	-	7	10	9	-	5
Death rate: Ischaemic heart disease	2	66	368	628	2,098	150
Death rate attributable to tobacco	-	5	38	54	-	9
Proportion of deaths attributable to tobacco (%)	-	8	10	9	-	6
Death rate: Cerebrovascular disease	-	17	80	192	644	41
Death rate attributable to tobacco	-	1	5	19	-	2
Proportion of deaths attributable to tobacco (%)	-	6	7	10	-	5
Death rate: Other cardiovascular diseases	50	155	575	1,301	6,680	374
Death rate attributable to tobacco	-	11	54	126	-	17
Proportion of deaths attributable to tobacco (%)	-	7	9	10	-	5
Death rate: Respiratory diseases	2	13	112	324	2,066	86
Death rate attributable to tobacco	-	1	22	92	-	8
Proportion of deaths attributable to tobacco (%)	-	4	20	28	-	9
Death rate: Chronic obstructive pulmonary disease	-	1	22	82	539	20
Death rate attributable to tobacco	-	0	11	54	-	5
Proportion of deaths attributable to tobacco (%)	-	15	53	66	-	22
Death rate: Other respiratory diseases	2	12	90	242	1,527	65
Death rate attributable to tobacco	-	0	11	38	-	4
Proportion of deaths attributable to tobacco (%)	-	3	12	16	-	5

Note 1: The total death rates are crude rates, and not standardized therefore cannot be used for comparison between countries/regions.

Note 2: The estimates above are based on SIR method (see Technical Note), some countries have used other methods which can produce somewhat different but equally valid estimates.

WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Cause of death	Men					Total for ages 30 & over	Women					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+		30-44	45-59	60-69	70-79	80+	
Death rate: ALL CAUSES	302	1,210	3,791	6,929	24,201	1,693	120	393	1,309	4,056	11,354	958
Death rate attributable to tobacco	-	64	577	896	-	124	-	-	-	328	-	23
Proportion of deaths attributable to tobacco (%)	-	5	15	13	-	7	-	-	-	8	-	2
Death rate: Communicable diseases	28	157	329	977	3,043	201	31	18	95	424	1,568	108
Death rate attributable to tobacco	-	2	9	37	-	3	-	-	-	21	-	1
Proportion of deaths attributable to tobacco (%)	-	1	3	4	-	2	-	-	-	5	-	1
Death rate: Tuberculosis	2	-	-	25	-	2	-	-	8	15	-	2
Death rate attributable to tobacco	-	-	-	3	-	0	-	-	-	1	-	0
Proportion of deaths attributable to tobacco (%)	-	-	-	12	-	7	-	-	-	8	-	4
Death rate: Lower respiratory infections	6	46	64	199	1,278	57	-	5	44	139	540	33
Death rate attributable to tobacco	-	2	9	33	-	3	-	-	-	19	-	1
Proportion of deaths attributable to tobacco (%)	-	4	14	17	-	6	-	-	-	14	-	4
Death rate: Noncommunicable diseases	160	856	3,249	5,842	21,158	1,342	76	325	1,206	3,618	9,787	825
Death rate attributable to tobacco	-	63	568	860	-	121	-	-	-	307	-	21
Proportion of deaths attributable to tobacco (%)	-	7	17	15	-	9	-	-	-	8	-	3
Death rate: All malignant neoplasms	28	198	806	1,307	1,884	259	25	171	285	642	169	149
Death rate attributable to tobacco	-	25	296	458	-	61	-	-	-	64	-	4
Proportion of deaths attributable to tobacco (%)	-	13	37	35	-	24	-	-	-	10	-	3
Death rate: Trachea, bronchus, lung cancers	-	12	103	241	-	27	-	5	8	76	-	8
Death rate attributable to tobacco	-	5	86	210	-	21	-	-	-	46	-	3
Proportion of deaths attributable to tobacco (%)	-	43	83	87	-	79	-	-	-	61	-	40
Death rate: All other malignant neoplasms	28	186	703	1,066	1,884	232	25	165	277	566	169	141
Death rate attributable to tobacco	-	20	210	248	-	40	-	-	-	18	-	1
Proportion of deaths attributable to tobacco (%)	-	11	30	23	-	17	-	-	-	3	-	1
Death rate: All Cardiovascular diseases	70	383	1,481	2,289	13,043	645	33	97	643	2,001	7,541	487
Death rate attributable to tobacco	-	35	216	216	-	44	-	-	-	187	-	13
Proportion of deaths attributable to tobacco (%)	-	9	15	9	-	7	-	-	-	9	-	3
Death rate: Ischaemic heart disease	4	125	614	834	3,483	205	-	9	163	481	1,379	96
Death rate attributable to tobacco	-	10	84	74	-	15	-	-	-	40	-	3
Proportion of deaths attributable to tobacco (%)	-	8	14	9	-	8	-	-	-	8	-	3
Death rate: Cerebrovascular disease	-	23	89	217	580	37	-	11	73	174	678	44
Death rate attributable to tobacco	-	2	12	17	-	3	-	-	-	20	-	1
Proportion of deaths attributable to tobacco (%)	-	9	13	8	-	7	-	-	-	11	-	3
Death rate: Other cardiovascular diseases	66	236	778	1,238	8,980	403	33	77	407	1,346	5,485	347
Death rate attributable to tobacco	-	23	120	125	-	26	-	-	-	127	-	9
Proportion of deaths attributable to tobacco (%)	-	10	15	10	-	6	-	-	-	9	-	3
Death rate: Respiratory diseases	2	21	207	539	4,492	131	2	5	33	170	805	43
Death rate attributable to tobacco	-	1	48	164	-	14	-	-	-	41	-	3
Proportion of deaths attributable to tobacco (%)	-	5	23	30	-	10	-	-	-	24	-	7
Death rate: Chronic obstructive pulmonary disease	-	1	39	145	1,179	32	-	1	7	37	206	10
Death rate attributable to tobacco	-	0	25	99	-	8	-	-	-	22	-	2
Proportion of deaths attributable to tobacco (%)	-	31	64	68	-	24	-	-	-	60	-	15
Death rate: Other respiratory diseases	2	20	167	394	3,314	100	2	4	27	133	599	33
Death rate attributable to tobacco	-	1	23	65	-	6	-	-	-	18	-	1
Proportion of deaths attributable to tobacco (%)	-	4	14	17	-	6	-	-	-	14	-	4

Note 3: For information on the sources of mortality data, please refer to Annex C, Table C4 of The global burden of disease: 2004 update.