


Sudan

Population 2004 (in thousands)

Age	Male	Female	Total
0-29	12,669	12,260	24,929
30-44	2,971	2,944	5,915
45-59	1,636	1,693	3,329
60-69	592	654	1,246
70-79	265	314	579
80+	62	85	147
Total	18,195	17,950	36,145

Population Pyramid



Highlights:

- The death rate from non-communicable diseases (1378 per 100,000 population) was about 3.2 times that for communicable diseases (434 per 100,000). Tobacco was responsible for 1% of all NCDs compared with 0% of all communicable disease related deaths.

- Within the non-communicable disease group, ischaemic heart disease accounted for 311 deaths per 100,000 population aged 30 years and over, with 1% of these deaths attributed to tobacco. Cancer of the trachea, bronchus and lung accounted for 6 deaths per 100,000 population but with 18% of these deaths attributed to tobacco.

- Within the communicable disease group, deaths attributed to tobacco accounted for 1% of all lower respiratory infection deaths and 0% of all tuberculosis deaths.

- The death rate due to tobacco in men was (21 per 100,000 men aged 30 years and over) compared with women (6 per 100,000 women aged 30 years and over). The proportion of deaths attributable to tobacco was almost 1% for men and 0% for women.

WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Cause of death	Both sexes					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+	
Death rate: ALL CAUSES	897	1,628	3,777	8,660	20,530	2,092
Death rate attributable to tobacco	-	10	45	85	84	13
Proportion of deaths attributable to tobacco (%)	-	1	1	1	0	1
Death rate: Communicable diseases	381	354	468	905	2,250	434
Death rate attributable to tobacco	-	0	2	4	8	1
Proportion of deaths attributable to tobacco (%)	-	0	0	0	0	0
Death rate: Tuberculosis	90	102	132	171	144	103
Death rate attributable to tobacco	-	0	1	1	1	0
Proportion of deaths attributable to tobacco (%)	-	0	1	1	0	0
Death rate: Lower respiratory infections	24	37	119	373	1,234	73
Death rate attributable to tobacco	-	0	1	3	7	0
Proportion of deaths attributable to tobacco (%)	-	0	1	1	1	1
Death rate: Noncommunicable diseases	221	1,028	3,050	7,474	17,692	1,378
Death rate attributable to tobacco	-	10	43	81	76	13
Proportion of deaths attributable to tobacco (%)	-	1	1	1	0	1
Death rate: All malignant neoplasms	55	204	448	791	1,088	194
Death rate attributable to tobacco	-	1	14	26	21	4
Proportion of deaths attributable to tobacco (%)	-	1	3	3	2	2
Death rate: Trachea, bronchus, lung cancers	1	7	18	22	24	6
Death rate attributable to tobacco	-	0	4	8	6	1
Proportion of deaths attributable to tobacco (%)	-	6	22	35	24	18
Death rate: All other malignant neoplasms	54	197	430	769	1,065	189
Death rate attributable to tobacco	-	1	10	18	15	3
Proportion of deaths attributable to tobacco (%)	-	1	2	2	1	1
Death rate: All Cardiovascular diseases	73	452	1,585	4,420	11,942	733
Death rate attributable to tobacco	-	7	13	17	4	4
Proportion of deaths attributable to tobacco (%)	-	1	1	0	0	1
Death rate: Ischaemic heart disease	15	199	771	1,948	4,381	311
Death rate attributable to tobacco	-	2	7	8	1	2
Proportion of deaths attributable to tobacco (%)	-	1	1	0	0	1
Death rate: Cerebrovascular disease	20	114	371	998	2,440	169
Death rate attributable to tobacco	-	2	3	3	1	1
Proportion of deaths attributable to tobacco (%)	-	2	1	0	0	1
Death rate: Other cardiovascular diseases	38	139	443	1,475	5,120	253
Death rate attributable to tobacco	-	3	4	6	2	2
Proportion of deaths attributable to tobacco (%)	-	2	1	0	0	1
Death rate: Respiratory diseases	15	64	226	599	1,399	101
Death rate attributable to tobacco	-	1	15	37	51	5
Proportion of deaths attributable to tobacco (%)	-	2	7	6	4	5
Death rate: Chronic obstructive pulmonary disease	4	34	142	397	874	60
Death rate attributable to tobacco	-	1	14	35	48	4
Proportion of deaths attributable to tobacco (%)	-	4	10	9	5	7
Death rate: Other respiratory diseases	10	30	84	202	524	41
Death rate attributable to tobacco	-	0	1	2	3	0
Proportion of deaths attributable to tobacco (%)	-	0	1	1	1	1

Note 1: The total death rates are crude rates, and not standardized therefore cannot be used for comparison between countries/regions.

Note 2: The estimates above are based on SIR method (see Technical Note), some countries have used other methods which can produce somewhat different but equally valid estimates.

WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Cause of death	Men					Total for ages 30 & over	Women					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+		30-44	45-59	60-69	70-79	80+	
Death rate: ALL CAUSES	1,148	2,010	4,561	9,719	22,015	2,414	643	1,258	3,068	7,768	19,449	1,779
Death rate attributable to tobacco	-	-	95	186	198	21	-	20	-	-	-	6
Proportion of deaths attributable to tobacco (%)	-	-	2	2	1	1	-	2	-	-	-	0
Death rate: Communicable diseases	413	468	637	1,078	2,623	510	350	243	314	760	1,979	361
Death rate attributable to tobacco	-	-	5	10	18	1	-	1	-	-	-	0
Proportion of deaths attributable to tobacco (%)	-	-	1	1	1	0	-	0	-	-	-	0
Death rate: Tuberculosis	133	156	189	249	228	152	46	50	80	105	83	55
Death rate attributable to tobacco	-	-	2	3	2	0	-	0	-	-	-	0
Proportion of deaths attributable to tobacco (%)	-	-	1	1	1	0	-	1	-	-	-	0
Death rate: Lower respiratory infections	34	54	163	451	1,563	91	14	21	80	307	995	54
Death rate attributable to tobacco	-	-	3	7	17	1	-	0	-	-	-	0
Proportion of deaths attributable to tobacco (%)	-	-	2	2	1	1	-	1	-	-	-	0
Death rate: Noncommunicable diseases	278	1,187	3,556	8,290	18,730	1,489	163	874	2,590	6,787	16,935	1,269
Death rate attributable to tobacco	-	-	90	176	180	20	-	19	-	-	-	6
Proportion of deaths attributable to tobacco (%)	-	-	3	2	1	1	-	2	-	-	-	0
Death rate: All malignant neoplasms	53	226	595	1,034	1,522	226	57	184	315	586	773	164
Death rate attributable to tobacco	-	-	29	56	49	6	-	3	-	-	-	1
Proportion of deaths attributable to tobacco (%)	-	-	5	5	3	3	-	2	-	-	-	1
Death rate: Trachea, bronchus, lung cancers	2	6	26	48	56	8	-	8	11	1	-	4
Death rate attributable to tobacco	-	-	8	17	14	2	-	1	-	-	-	0
Proportion of deaths attributable to tobacco (%)	-	-	33	35	24	22	-	11	-	-	-	7
Death rate: All other malignant neoplasms	51	220	570	986	1,466	217	57	176	304	586	773	161
Death rate attributable to tobacco	-	-	21	39	35	5	-	2	-	-	-	1
Proportion of deaths attributable to tobacco (%)	-	-	4	4	2	2	-	1	-	-	-	0
Death rate: All Cardiovascular diseases	95	529	1,790	4,692	12,036	759	50	377	1,400	4,191	11,873	708
Death rate attributable to tobacco	-	-	28	37	8	5	-	13	-	-	-	4
Proportion of deaths attributable to tobacco (%)	-	-	2	1	0	1	-	4	-	-	-	1
Death rate: Ischaemic heart disease	24	267	938	2,290	4,614	354	6	134	621	1,659	4,211	269
Death rate attributable to tobacco	-	-	14	18	3	2	-	4	-	-	-	1
Proportion of deaths attributable to tobacco (%)	-	-	2	1	0	1	-	3	-	-	-	0
Death rate: Cerebrovascular disease	25	119	400	1,012	2,359	166	14	108	346	985	2,500	171
Death rate attributable to tobacco	-	-	6	7	2	1	-	4	-	-	-	1
Proportion of deaths attributable to tobacco (%)	-	-	1	1	0	1	-	4	-	-	-	1
Death rate: Other cardiovascular diseases	46	143	453	1,390	5,063	239	29	134	434	1,547	5,162	267
Death rate attributable to tobacco	-	-	8	13	4	2	-	5	-	-	-	1
Proportion of deaths attributable to tobacco (%)	-	-	2	1	0	1	-	4	-	-	-	1
Death rate: Respiratory diseases	21	88	312	773	1,716	127	9	42	148	453	1,167	76
Death rate attributable to tobacco	-	-	31	81	120	9	-	3	-	-	-	1
Proportion of deaths attributable to tobacco (%)	-	-	10	10	7	7	-	6	-	-	-	1
Death rate: Chronic obstructive pulmonary disease	7	47	202	524	1,075	77	2	22	88	290	728	44
Death rate attributable to tobacco	-	-	30	77	113	8	-	2	-	-	-	1
Proportion of deaths attributable to tobacco (%)	-	-	15	15	11	11	-	11	-	-	-	2
Death rate: Other respiratory diseases	14	41	110	249	642	50	7	20	60	163	439	32
Death rate attributable to tobacco	-	-	2	4	7	0	-	0	-	-	-	0
Proportion of deaths attributable to tobacco (%)	-	-	2	2	1	1	-	1	-	-	-	0

Note 3: For information on the sources of mortality data, please refer to Annex C, Table C4 of The global burden of disease: 2004 update.