


Togo

Population 2004 (in thousands)

Age	Male	Female	Total
0-29	2,170	2,177	4,348
30-44	460	470	931
45-59	242	261	503
60-69	84	98	182
70-79	38	50	87
80+	8	12	20
Total	3,003	3,068	6,071

Population Pyramid



Highlights:

- The death rate from non-communicable diseases (1090 per 100,000 population) was about 1.4 times that for communicable diseases (766 per 100,000). Tobacco was responsible for 3% of all NCDs compared with 1% of all communicable disease related deaths.

- Within the non-communicable disease group, ischaemic heart disease accounted for 152 deaths per 100,000 population aged 30 years and over, with 2% of these deaths attributed to tobacco. Cancer of the trachea, bronchus and lung accounted for 8 deaths per 100,000 population but with 41% of these deaths attributed to tobacco.

- Within the communicable disease group, deaths attributed to tobacco accounted for 2% of all lower respiratory infection deaths and 1% of all tuberculosis deaths.

- The death rate due to tobacco in men was (68 per 100,000 men aged 30 years and over) compared with women (0 per 100,000 women aged 30 years and over). The proportion of deaths attributable to tobacco was almost 3% for men and 0% for women.

WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Cause of death	Both sexes					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+	
Death rate: ALL CAUSES	934	1,649	3,510	7,890	17,802	1,961
Death rate attributable to tobacco	-	43	94	147	287	33
Proportion of deaths attributable to tobacco (%)	-	3	3	2	2	2
Death rate: Communicable diseases	623	737	975	1,542	2,908	766
Death rate attributable to tobacco	-	7	13	18	46	5
Proportion of deaths attributable to tobacco (%)	-	1	1	1	2	1
Death rate: Tuberculosis	143	169	226	282	265	168
Death rate attributable to tobacco	-	4	5	5	6	2
Proportion of deaths attributable to tobacco (%)	-	2	2	2	2	1
Death rate: Lower respiratory infections	6	99	331	697	1,621	121
Death rate attributable to tobacco	-	3	9	13	39	3
Proportion of deaths attributable to tobacco (%)	-	3	3	2	2	2
Death rate: Noncommunicable diseases	223	811	2,404	6,139	14,599	1,090
Death rate attributable to tobacco	-	36	80	129	241	28
Proportion of deaths attributable to tobacco (%)	-	4	3	2	2	3
Death rate: All malignant neoplasms	35	175	494	1,106	1,981	201
Death rate attributable to tobacco	-	8	21	33	63	7
Proportion of deaths attributable to tobacco (%)	-	4	4	3	3	3
Death rate: Trachea, bronchus, lung cancers	1	7	23	41	56	8
Death rate attributable to tobacco	-	3	11	17	27	3
Proportion of deaths attributable to tobacco (%)	-	40	48	41	48	41
Death rate: All other malignant neoplasms	34	168	471	1,065	1,925	193
Death rate attributable to tobacco	-	5	11	16	37	4
Proportion of deaths attributable to tobacco (%)	-	3	2	2	2	2
Death rate: All Cardiovascular diseases	74	328	1,062	3,028	8,140	495
Death rate attributable to tobacco	-	18	22	23	11	9
Proportion of deaths attributable to tobacco (%)	-	5	2	1	0	2
Death rate: Ischaemic heart disease	14	110	388	942	2,049	152
Death rate attributable to tobacco	-	7	9	9	3	3
Proportion of deaths attributable to tobacco (%)	-	6	2	1	0	2
Death rate: Cerebrovascular disease	30	123	392	1,121	2,883	183
Death rate attributable to tobacco	-	6	7	7	3	3
Proportion of deaths attributable to tobacco (%)	-	5	2	1	0	2
Death rate: Other cardiovascular diseases	30	96	282	965	3,208	160
Death rate attributable to tobacco	-	5	6	8	5	2
Proportion of deaths attributable to tobacco (%)	-	5	2	1	0	2
Death rate: Respiratory diseases	29	91	281	662	1,384	121
Death rate attributable to tobacco	-	10	35	70	160	12
Proportion of deaths attributable to tobacco (%)	-	11	13	11	12	10
Death rate: Chronic obstructive pulmonary disease	6	36	139	339	646	53
Death rate attributable to tobacco	-	9	32	64	143	11
Proportion of deaths attributable to tobacco (%)	-	24	23	19	22	20
Death rate: Other respiratory diseases	23	56	142	323	738	69
Death rate attributable to tobacco	-	1	3	6	18	1
Proportion of deaths attributable to tobacco (%)	-	3	2	2	2	2

Note 1: The total death rates are crude rates, and not standardized therefore cannot be used for comparison between countries/regions.

Note 2: The estimates above are based on SIR method (see Technical Note), some countries have used other methods which can produce somewhat different but equally valid estimates.

WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Cause of death	Men					Total for ages 30 & over	Women					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+		30-44	45-59	60-69	70-79	80+	
Death rate: ALL CAUSES	957	1,949	4,118	8,748	19,936	2,094	912	1,369	2,987	7,238	16,455	1,838
Death rate attributable to tobacco	-	88	203	341	742	68	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	5	5	4	4	3	-	-	-	-	-	-
Death rate: Communicable diseases	588	891	1,253	1,896	3,550	830	657	593	736	1,273	2,502	707
Death rate attributable to tobacco	-	14	29	42	119	10	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	2	2	2	3	1	-	-	-	-	-	-
Death rate: Tuberculosis	219	266	334	435	458	257	68	79	133	165	144	85
Death rate attributable to tobacco	-	8	10	11	17	4	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	3	3	2	4	2	-	-	-	-	-	-
Death rate: Lower respiratory infections	6	115	443	858	1,962	138	7	85	234	574	1,406	105
Death rate attributable to tobacco	-	5	19	31	101	6	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	5	4	4	5	4	-	-	-	-	-	-
Death rate: Noncommunicable diseases	230	899	2,683	6,552	16,036	1,105	216	729	2,163	5,825	13,692	1,076
Death rate attributable to tobacco	-	75	174	299	623	59	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	8	6	5	4	5	-	-	-	-	-	-
Death rate: All malignant neoplasms	28	181	578	1,374	2,866	215	42	170	421	903	1,422	188
Death rate attributable to tobacco	-	16	46	77	163	14	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	9	8	6	6	7	-	-	-	-	-	-
Death rate: Trachea, bronchus, lung cancers	1	13	41	70	111	13	0	2	7	19	21	3
Death rate attributable to tobacco	-	6	23	39	69	7	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	46	57	56	62	51	-	-	-	-	-	-
Death rate: All other malignant neoplasms	27	168	538	1,303	2,755	203	41	169	414	884	1,402	185
Death rate attributable to tobacco	-	10	23	38	94	8	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	6	4	3	3	4	-	-	-	-	-	-
Death rate: All Cardiovascular diseases	72	354	1,113	2,910	8,027	461	76	304	1,018	3,117	8,211	526
Death rate attributable to tobacco	-	36	48	54	29	18	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	10	4	2	0	4	-	-	-	-	-	-
Death rate: Ischaemic heart disease	17	148	483	1,159	2,425	177	10	74	305	778	1,812	128
Death rate attributable to tobacco	-	14	20	21	8	7	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	10	4	2	0	4	-	-	-	-	-	-
Death rate: Cerebrovascular disease	28	125	382	966	2,530	158	32	120	401	1,239	3,105	207
Death rate attributable to tobacco	-	13	15	15	8	6	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	10	4	2	0	4	-	-	-	-	-	-
Death rate: Other cardiovascular diseases	27	80	248	785	3,072	127	34	110	312	1,101	3,294	190
Death rate attributable to tobacco	-	10	13	18	13	5	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	12	5	2	0	4	-	-	-	-	-	-
Death rate: Respiratory diseases	30	121	392	912	1,881	150	28	64	185	472	1,069	94
Death rate attributable to tobacco	-	21	76	162	415	25	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	17	20	18	22	17	-	-	-	-	-	-
Death rate: Chronic obstructive pulmonary disease	7	52	214	512	991	73	5	20	74	208	428	34
Death rate attributable to tobacco	-	18	69	148	369	22	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	34	32	29	37	30	-	-	-	-	-	-
Death rate: Other respiratory diseases	23	69	178	400	890	77	23	44	111	264	641	61
Death rate attributable to tobacco	-	3	7	14	46	3	-	-	-	-	-	-
Proportion of deaths attributable to tobacco (%)	-	5	4	4	5	4	-	-	-	-	-	-

Note 3: For information on the sources of mortality data, please refer to Annex C, Table C4 of The global burden of disease: 2004 update.