


## Uganda

### Population 2004 (in thousands)

Age	Male	Female	Total
0-29	10,836	10,705	21,541
30-44	1,814	1,754	3,568
45-59	870	968	1,838
60-69	292	350	642
70-79	155	196	351
80+	37	52	88
<b>Total</b>	<b>14,004</b>	<b>14,024</b>	<b>28,028</b>

### Population Pyramid



### Highlights:

- The death rate from non-communicable diseases (1075 per 100,000 population) was about 0.8 times that for communicable diseases (1423 per 100,000). Tobacco was responsible for 2% of all NCDs compared with 0% of all communicable disease related deaths.

- Within the non-communicable disease group, ischaemic heart disease accounted for 153 deaths per 100,000 population aged 30 years and over, with 2% of these deaths attributed to tobacco. Cancer of the trachea, bronchus and lung accounted for 8 deaths per 100,000 population but with 26% of these deaths attributed to tobacco.

- Within the communicable disease group, deaths attributed to tobacco accounted for 1% of all lower respiratory infection deaths and 1% of all tuberculosis deaths.

- The death rate due to tobacco in men was (46 per 100,000 men aged 30 years and over) compared with women (13 per 100,000 women aged 30 years and over). The proportion of deaths attributable to tobacco was almost 2% for men and 0% for women.

## WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Cause of death	Both sexes					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+	
<b>Death rate: ALL CAUSES</b>	<b>1,788</b>	<b>2,400</b>	<b>3,772</b>	<b>7,982</b>	<b>18,030</b>	<b>2,714</b>
Death rate attributable to tobacco	-	46	89	122	56	29
Proportion of deaths attributable to tobacco (%)	-	2	2	2	0	1
<b>Death rate: Communicable diseases</b>	<b>1,442</b>	<b>1,370</b>	<b>1,101</b>	<b>1,625</b>	<b>3,256</b>	<b>1,423</b>
Death rate attributable to tobacco	-	5	9	14	9	3
Proportion of deaths attributable to tobacco (%)	-	0	1	1	0	0
<b>Death rate: Tuberculosis</b>	<b>149</b>	<b>219</b>	<b>256</b>	<b>257</b>	<b>231</b>	<b>186</b>
Death rate attributable to tobacco	-	5	3	2	1	2
Proportion of deaths attributable to tobacco (%)	-	2	1	1	0	1
<b>Death rate: Lower respiratory infections</b>	<b>8</b>	<b>39</b>	<b>244</b>	<b>697</b>	<b>1,855</b>	<b>103</b>
Death rate attributable to tobacco	-	1	6	12	8	2
Proportion of deaths attributable to tobacco (%)	-	2	2	2	0	1
<b>Death rate: Noncommunicable diseases</b>	<b>146</b>	<b>834</b>	<b>2,413</b>	<b>6,025</b>	<b>14,245</b>	<b>1,075</b>
Death rate attributable to tobacco	-	40	80	108	47	26
Proportion of deaths attributable to tobacco (%)	-	5	3	2	0	2
<b>Death rate: All malignant neoplasms</b>	<b>21</b>	<b>181</b>	<b>496</b>	<b>1,091</b>	<b>1,932</b>	<b>197</b>
Death rate attributable to tobacco	-	12	26	26	11	8
Proportion of deaths attributable to tobacco (%)	-	7	5	2	1	4
<b>Death rate: Trachea, bronchus, lung cancers</b>	<b>1</b>	<b>9</b>	<b>26</b>	<b>37</b>	<b>44</b>	<b>8</b>
Death rate attributable to tobacco	-	3	9	6	2	2
Proportion of deaths attributable to tobacco (%)	-	33	36	17	5	26
<b>Death rate: All other malignant neoplasms</b>	<b>20</b>	<b>172</b>	<b>470</b>	<b>1,054</b>	<b>1,888</b>	<b>189</b>
Death rate attributable to tobacco	-	9	17	20	9	5
Proportion of deaths attributable to tobacco (%)	-	5	4	2	0	3
<b>Death rate: All Cardiovascular diseases</b>	<b>47</b>	<b>338</b>	<b>1,064</b>	<b>2,963</b>	<b>7,945</b>	<b>496</b>
Death rate attributable to tobacco	-	18	20	42	8	9
Proportion of deaths attributable to tobacco (%)	-	5	2	1	0	2
<b>Death rate: Ischaemic heart disease</b>	<b>10</b>	<b>113</b>	<b>389</b>	<b>927</b>	<b>2,005</b>	<b>153</b>
Death rate attributable to tobacco	-	7	8	8	2	3
Proportion of deaths attributable to tobacco (%)	-	6	2	1	0	2
<b>Death rate: Cerebrovascular disease</b>	<b>19</b>	<b>125</b>	<b>390</b>	<b>1,087</b>	<b>2,780</b>	<b>181</b>
Death rate attributable to tobacco	-	6	6	18	0	3
Proportion of deaths attributable to tobacco (%)	-	5	2	2	0	2
<b>Death rate: Other cardiovascular diseases</b>	<b>19</b>	<b>99</b>	<b>285</b>	<b>949</b>	<b>3,160</b>	<b>161</b>
Death rate attributable to tobacco	-	5	5	15	5	3
Proportion of deaths attributable to tobacco (%)	-	5	2	2	0	2
<b>Death rate: Respiratory diseases</b>	<b>19</b>	<b>94</b>	<b>282</b>	<b>655</b>	<b>1,356</b>	<b>119</b>
Death rate attributable to tobacco	-	10	33	37	27	8
Proportion of deaths attributable to tobacco (%)	-	11	12	6	2	7
<b>Death rate: Chronic obstructive pulmonary disease</b>	<b>4</b>	<b>37</b>	<b>140</b>	<b>337</b>	<b>634</b>	<b>53</b>
Death rate attributable to tobacco	-	9	30	32	24	7
Proportion of deaths attributable to tobacco (%)	-	24	21	9	4	14
<b>Death rate: Other respiratory diseases</b>	<b>15</b>	<b>57</b>	<b>143</b>	<b>318</b>	<b>722</b>	<b>66</b>
Death rate attributable to tobacco	-	1	3	5	3	1
Proportion of deaths attributable to tobacco (%)	-	3	2	2	0	2

Note 1: The total death rates are crude rates, and not standardized therefore cannot be used for comparison between countries/regions.

Note 2: The estimates above are based on SIR method (see Technical Note), some countries have used other methods which can produce somewhat different but equally valid estimates.

## WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Cause of death	Men					Total for ages 30 & over	Women					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+		30-44	45-59	60-69	70-79	80+	
<b>Death rate: ALL CAUSES</b>	1,732	2,759	4,466	8,821	19,377	2,817	1,846	2,078	3,193	7,321	17,069	2,616
Death rate attributable to tobacco	-	96	196	19	41	46	-	-	-	203	67	13
Proportion of deaths attributable to tobacco (%)	-	3	4	0	0	2	-	-	-	3	0	0
<b>Death rate: Communicable diseases</b>	1,191	1,520	1,344	1,890	3,728	1,359	1,702	1,236	897	1,416	2,919	1,483
Death rate attributable to tobacco	-	11	20	2	6	5	-	-	-	23	11	2
Proportion of deaths attributable to tobacco (%)	-	1	2	0	0	0	-	-	-	2	0	0
<b>Death rate: Tuberculosis</b>	205	320	291	350	333	253	91	128	227	183	158	123
Death rate attributable to tobacco	-	10	8	0	1	3	-	-	-	3	1	0
Proportion of deaths attributable to tobacco (%)	-	3	3	0	0	1	-	-	-	2	0	0
<b>Death rate: Lower respiratory infections</b>	1	34	338	834	2,152	107	16	43	165	589	1,643	99
Death rate attributable to tobacco	-	1	13	1	5	2	-	-	-	20	11	1
Proportion of deaths attributable to tobacco (%)	-	4	4	0	0	2	-	-	-	3	1	1
<b>Death rate: Noncommunicable diseases</b>	212	938	2,755	6,504	15,079	1,125	78	741	2,126	5,648	13,651	1,028
Death rate attributable to tobacco	-	85	176	17	36	41	-	-	-	180	55	11
Proportion of deaths attributable to tobacco (%)	-	9	6	0	0	4	-	-	-	3	0	1
<b>Death rate: All malignant neoplasms</b>	26	189	594	1,364	2,687	219	16	173	414	876	1,393	176
Death rate attributable to tobacco	-	25	58	5	9	13	-	-	-	43	13	3
Proportion of deaths attributable to tobacco (%)	-	13	10	0	0	6	-	-	-	5	1	2
<b>Death rate: Trachea, bronchus, lung cancers</b>	2	13	38	33	45	10	1	5	16	40	42	6
Death rate attributable to tobacco	-	6	21	2	3	4	-	-	-	10	2	1
Proportion of deaths attributable to tobacco (%)	-	46	54	5	7	36	-	-	-	25	5	10
<b>Death rate: All other malignant neoplasms</b>	24	176	556	1,331	2,642	209	15	168	399	836	1,351	170
Death rate attributable to tobacco	-	19	37	3	6	9	-	-	-	33	11	2
Proportion of deaths attributable to tobacco (%)	-	11	7	0	0	4	-	-	-	4	1	1
<b>Death rate: All Cardiovascular diseases</b>	66	369	1,143	2,889	7,555	474	28	309	999	3,021	8,223	517
Death rate attributable to tobacco	-	37	44	2	1	14	-	-	-	73	12	4
Proportion of deaths attributable to tobacco (%)	-	10	4	0	0	3	-	-	-	2	0	1
<b>Death rate: Ischaemic heart disease</b>	16	155	497	1,152	2,285	180	4	75	299	750	1,805	128
Death rate attributable to tobacco	-	15	18	1	0	6	-	-	-	14	4	1
Proportion of deaths attributable to tobacco (%)	-	9	4	0	0	3	-	-	-	2	0	1
<b>Death rate: Cerebrovascular disease</b>	26	131	393	961	2,387	162	12	120	388	1,186	3,060	200
Death rate attributable to tobacco	-	13	14	1	0	5	-	-	-	32	-	2
Proportion of deaths attributable to tobacco (%)	-	10	4	0	0	3	-	-	-	3	-	1
<b>Death rate: Other cardiovascular diseases</b>	25	83	253	776	2,883	132	12	114	312	1,085	3,358	189
Death rate attributable to tobacco	-	10	12	1	1	4	-	-	-	26	9	2
Proportion of deaths attributable to tobacco (%)	-	12	5	0	0	3	-	-	-	2	0	1
<b>Death rate: Respiratory diseases</b>	28	126	402	905	1,767	153	10	65	182	458	1,062	87
Death rate attributable to tobacco	-	21	72	10	25	13	-	-	-	58	28	4
Proportion of deaths attributable to tobacco (%)	-	17	18	1	1	9	-	-	-	13	3	4
<b>Death rate: Chronic obstructive pulmonary disease</b>	7	55	219	509	930	75	2	20	73	201	423	33
Death rate attributable to tobacco	-	18	65	9	23	12	-	-	-	50	24	3
Proportion of deaths attributable to tobacco (%)	-	33	30	2	2	16	-	-	-	25	6	10
<b>Death rate: Other respiratory diseases</b>	21	72	183	397	837	78	8	44	109	257	639	54
Death rate attributable to tobacco	-	3	7	1	2	2	-	-	-	9	4	1
Proportion of deaths attributable to tobacco (%)	-	4	4	0	0	2	-	-	-	3	1	1

Note 3: For information on the sources of mortality data, please refer to Annex C, Table C4 of The global burden of disease: 2004 update.